arXiv:1412.0319v1 [math.SP] 1 Dec 2014

Spectra of the blow-up graphs

Carla Oliveira*† Leonardo de Lima‡§ and Vladimir Nikiforov¶

Abstract

Let G be graph on n vertices and $G^{(t)}$ its blow-up graph of order t. In this paper, we determine all eigenvalues of the Laplacian and the signless Laplacian matrix of $G^{(t)}$ and its complement $\overline{G^{(t)}}$.

Keywords: blow-up graphs, eigenvalues, Laplacian, signless Laplacian.

1 Introduction

Let G be a graph on n vertices with degree sequence $d_1, d_2, \ldots, d_n$ in non-incresing order. We denote the complement of G by $\overline{G}$. Write $A = A(G)$ for the adjacency matrix of G and let $D = D(G)$ be the diagonal matrix of the row-sums of A, i.e., the degrees of G. The Laplacian $L(G)$ and signless Laplacian $Q(G)$ of G are defined as $L(G) = D - A$ and $Q(G) = D + A$. The eigenvalues of $A(G)$, $L(G)$ and $Q(G)$ arranged in non-increasing order are denoted by $\lambda_1, \ldots, \lambda_n$, $\mu_1, \ldots, \mu_n$ and $q_1, \ldots, q_n$.

For any graph G and integer $t \geq 1$, write $G^{(t)}$ for the graph obtained by replacing each vertex u of G by a set V_u of t independent vertices and every edge $\{u, v\}$ of G by a complete bipartite graph with parts V_u and V_v. Usually $G^{(t)}$ is called a *blow-up* of G. See Figure 1 for a blow-up example of order 2 and 3.

The blow-up of a graph G has a useful algebraic characterization: if A is the adjacency matrix of G, then the adjacency matrix $A\left(G^{(t)}\right)$ of $G^{(t)}$ is given by

$$A\left(G^{(t)}\right) = A \otimes J_t$$

where $\otimes$ is the Kronecker product and J_t is the all ones matrix of order t. This observation yields the following fact.

*National School of Statistics, Rio de Janeiro, Brazil;*email: carla.oliveira@ibge.gov.br*

†Research supported by CNPq Grant 305454/2012-9.

‡Department of Production Engineering, Federal Center of Technological Education, Rio de Janeiro, Brazil;*email: llima@cefet-rj.br*

§Research supported by Programa Jovem Cientista do Nosso Estado, FAPERJ Grant E-26/102.218/2013, and CNPq Grant 305867/2012-1.

¶Department of Mathematical Sciences, University of Memphis, Memphis TN 38152, USA; *email: vnikifrv@memphis.edu*



Figure 1: A graph G and its blow-up of order 2 and 3.

Proposition 1 *The eigenvalues of* $G^{(t)}$ *are* $t\lambda_1(G), \ldots, t\lambda_n(G)$, *together with* $n(t-1)$ *additional* 0*'s.*

For the complements of graph blow-ups, one can easily check the following fact.

Proposition 2 *The eigenvalues of* $\overline{G^{(t)}}$ *are* $t\lambda_1\left(\overline{G}\right)+t-1, \ldots, t\lambda_n\left(\overline{G}\right)+t-1$, *together with* $n(t-1)$ *additional* (-1)*'s.*

The goal of this paper is to determine the spectra of $L(G^{(t)})$, $Q(G^{(t)})$, and $Q(\overline{G^{(t)}})$, which is a more difficult task than for the adjacency matrix. We also note that the spectrum of $L(\overline{G^{(t)}})$ is obtained immediately from our results as $\mu_{n-i}(\overline{G}) = n - \mu_i(G)$, for $i = 1, \ldots, n-1$, (see e.g. [1]).

2 Main Results

In this section, we prove the main results of the paper.

Theorem 3 *If* G *is a graph on* n *vertices, the Laplacian eigenvalues of* $G^{(t)}$ *are* $t\mu_1, \ldots, t\mu_n$ *and* $td_1, \ldots, td_n$, *where each* td_i *has multiplicity* $(t-1)$ *for* $i = 1, \ldots, n$.

Proof The Laplacian matrix of $G^{(t)}$ can be written as

$$L(G^{(t)}) = \begin{bmatrix} L(G)+(t-1)D & -A(G) & \ldots & -A(G) \\ -A(G) & L(G)+(t-1)D & \ldots & -A(G) \\ -A(G) & -A(G) & \ldots & -A(G) \\ \vdots & & \ddots & \vdots \\ -A(G) & -A(G) & \ldots & L(G)+(t-1)D \end{bmatrix},$$

where $L(G)+(t-1)D$ is the matrix of order n and D is the diagonal matrix of the vertices degree. Let $\{\mathbf{x}_i\}$ be an orthogonal basis of eigenvectors to $L(G)$, such that for each $i = 1, \ldots, n$, $\mathbf{x}_i$ is an

eigenvector to μ_i. Consider the $tn-$vector $\mathbf{y}_i$ such that ${\mathbf{y}_i}^T = [\mathbf{x}_i \ \ \mathbf{x}_i \ \ \cdots \ \ \mathbf{x}_i]$, for $i = 1, \ldots, n$. Observe that

$$L(G^{(t)})\mathbf{y}_i = \begin{bmatrix} L(G)\mathbf{x}_i + (t-1)D\mathbf{x}_i - (t-1)A(G)\mathbf{x}_i \\ \hline L(G)\mathbf{x}_i + (t-1)D\mathbf{x}_i - (t-1)A(G)\mathbf{x}_i \\ \hline \vdots \\ \hline L(G)\mathbf{x}_i + (t-1)D\mathbf{x}_i - (t-1)A(G)\mathbf{x}_i \end{bmatrix} = \begin{bmatrix} tL(G)\mathbf{x}_i \\ \hline tL(G)\mathbf{x}_i \\ \hline \vdots \\ \hline tL(G)\mathbf{x}_i \end{bmatrix} = \begin{bmatrix} t\mu_i\mathbf{x}_i \\ \hline t\mu_i\mathbf{x}_i \\ \hline \vdots \\ \hline t\mu_i\mathbf{x}_i \end{bmatrix} = t\mu_i\mathbf{y}_i.$$

So $\mathbf{y}_i$ is an eigenvector to $t\mu_i$. It easy to see that $\mathbf{y}_i^T\mathbf{y}_j = 0$, since $\mathbf{x}_i^T\mathbf{x}_j = 0$, for all $i \neq j$, $1 \leq i, j \leq n$. Consider the tn-vector E_i of the form ${E_i}^T = [\alpha_{i1} \ \ \alpha_{i2} \ \ \cdots \ \ \alpha_{it}]$ such that, $\alpha_{ik} \in \Re^n$ for all $1 \leq k \leq t$. For each $i = 1, \ldots, n$, let e_i the $i-th$ standard unit basis vector. For each $k = 1, \ldots, t-1$, we define the nt-vector

$$({E_i}^k)^T = [0 \ \ \cdots \ \ 0 \ \ e_{ik} \ \ -e_{i(k+1)} \ \ 0 \ \ \cdots \ \ 0]$$

and we get $L(G^{(t)}){E_i}^k = td_i{E_i}^k$. So ${E_i}^1, {E_i}^2, \ldots, {E_i}^{t-1}$ are eigenvectors to td_i. Now, we need to prove that the set $\{{E_i}^1, {E_i}^2, \ldots, {E_i}^{t-1}\}$ is linearly independent. Suppose that $\sum_{j=1}^{t-1} c_j{E_i}^j = 0$. So $c_1 = c_{t-1} = 0$ and $-c_j + c_{j+1} = 0$, for all $j = 1, \ldots, t-2$. Then $c_j = 0$, $1 \leq j \leq t-1$ which implies that td_i is an eigenvalue of multiplicity at least $t-1$. Also, $\mathbf{x}_j^T E_i^k = 0$ for $i \neq j$, $1 \leq i, j \leq n$ and $1 \leq k \leq t-1$ and the result follows. □

Theorem 4 *If G be is a graph on n vertices, the signless Laplacian eigenvalues of $G^{(t)}$ are $tq_1, \ldots, tq_n$ and $td_1, \ldots, td_n$, where each td_i has multiplicity $(t-1)$ for $i = 1, \ldots, n$.*

Proof The signless Laplacian matrix of $G^{(t)}$ can be written as

$$Q(G^{(t)}) = \begin{bmatrix} Q(G) + (t-1)D & A(G) & \ldots & A(G) \\ A(G) & Q(G) + (t-1)D & \ldots & A(G) \\ A(G) & A(G) & \ldots & A(G) \\ \vdots & & \ddots & \vdots \\ A(G) & A(G) & \ldots & Q(G) + (t-1)D \end{bmatrix},$$

where $Q(G) + (t-1)D$ is the matrix of order n and D is the diagonal matrix of the vertices degree. The proof follows analogously to Theorem 3 taking the same eigenvectors ${\mathbf{y}_i}^T = [\mathbf{x}_i \ \ \mathbf{x}_i \ \ \cdots \ \ \mathbf{x}_i]$, $\mathbf{y}_i$ is an eigenvector to q_i for $i = 1, \ldots, n$, and also

$$({E_i}^k)^T = [0 \ \ \cdots \ \ 0 \ \ e_{ik} \ \ -e_{i(k+1)} \ \ 0 \ \ \cdots \ \ 0]$$

for each $i = 1, \ldots, n$ and $k = 1, \ldots, t-1$, and the result follows. □

Theorem 5 *If G is a graph on n vertices, the signless Laplacian eigenvalues of $\overline{G^{(t)}}$ are $t\overline{q}_1 + 2(t-1), \ldots, t\overline{q}_n + 2(t-1)$ and $tn - td_1 - 2, \ldots, tn - td_n - 2$, where each $tn - td_i - 2$ has multiplicity $(t-1)$ for $i = 1, \ldots, n$.*

Proof With a convenient labeling of the blow-up graph $G^{(t)}$, we can write the signless Laplacian of the complement of $G^{(t)}$ as the following

$$Q(\overline{G^{(t)}}) = \begin{bmatrix} (tn-2)I_n - tD + J_n - A & \cdots & J_n - A(G) \\ J_n - A(G) & \ddots & J_n - A(G) \\ \vdots & \vdots & \vdots \\ J_n - A(G) & \cdots & (tn-2)I_n - tD + J_n - A \end{bmatrix}.$$

Considering the $tn-$vectors ${E_i}^k$ for each $k = 1, \ldots, n$ we get

$$Q(\overline{G^{(t)}}){E_i}^k = \begin{pmatrix} 0 & \ldots & (tn-2-td_i) & \ldots & 0 & 0 & \ldots & -(tn-2-td_i) & \ldots & 0 & 0 & \ldots & 0 \end{pmatrix}^T$$

and then $Q(\overline{G^{(t)}}){E_i}^k = (tn-2-td_i){E_i}^k$. So ${E_i}^1, {E_i}^2, \ldots, {E_i}^{t-1}$ are eigenvectors to $tn-2-td_i$ for all $i = 1, \ldots, n$. Since the set $\{{E_i}^1, {E_i}^2, \ldots, {E_i}^{t-1}\}$ is linearly independent, the eigenvalue $tn-2-td_i$ has multiplicity at least $t-1$. Therefore, $nt-n$ eigenvalues of $Q(\overline{G^{(t)}})$ are known and we need to find the remaining n. One can easily rewrite $Q(\overline{G^{(t)}})$ in a such way that the block matrix $Q(\overline{G}) + (t-1)nI_n - (t-1)D$ appears in the main diagonal and $A(\overline{G}) + I_n$ in the remaining positions. Let $\{\mathbf{x}_i\}$ be an orthogonal basis of eigenvectors to $Q(\overline{G})$, such that for each $i = 1, \ldots, n$, $\mathbf{x}_i$ is an eigenvector to $\overline{q}_i$. Consider the $tn-$vectors $\mathbf{y}_i$ such that ${\mathbf{y}_i}^T = [\mathbf{x}_i \ \ \mathbf{x}_i \ \ \cdots \ \ \mathbf{x}_i]$, for $i = 1, \ldots, n$. Observe that

$$Q(\overline{G^{(t)}})\mathbf{y}_i = \begin{bmatrix} Q(\overline{G})\mathbf{x}_i + (t-1)n\mathbf{x}_i - (t-1)D\mathbf{x}_i + (t-1)(A(\overline{G}) + I_n)\mathbf{x}_i \\ \hline Q(\overline{G})\mathbf{x}_i + (t-1)n\mathbf{x}_i - (t-1)D\mathbf{x}_i + (t-1)(A(\overline{G}) + I_n)\mathbf{x}_i \\ \hline \vdots \\ \hline Q(\overline{G})\mathbf{x}_i + (t-1)n\mathbf{x}_i - (t-1)D\mathbf{x}_i + (t-1)(A(\overline{G}) + I_n)\mathbf{x}_i \end{bmatrix} =$$

$$= \begin{bmatrix} \overline{q}_i\mathbf{x}_i + (t-1)(n-1)I_n\mathbf{x}_i - (t-1)D\mathbf{x}_i + (t-1)A(\overline{G})\mathbf{x}_i + 2(t-1)\mathbf{x}_i \\ \hline \overline{q}_i\mathbf{x}_i + (t-1)(n-1)I_n\mathbf{x}_i - (t-1)D\mathbf{x}_i + (t-1)A(\overline{G})\mathbf{x}_i + 2(t-1)\mathbf{x}_i \\ \hline \vdots \\ \hline \overline{q}_i\mathbf{x}_i + (t-1)(n-1)I_n\mathbf{x}_i - (t-1)D\mathbf{x}_i + (t-1)A(\overline{G})\mathbf{x}_i + 2(t-1)\mathbf{x}_i \end{bmatrix} =$$

$$= \begin{bmatrix} \overline{q}_i\mathbf{x}_i + ((t-1)(n-1)I_n - (t-1)D)\mathbf{x}_i + (t-1)A(\overline{G})\mathbf{x}_i + 2(t-1)\mathbf{x}_i \\ \hline \overline{q}_i\mathbf{x}_i + ((t-1)(n-1)I_n - (t-1)D)\mathbf{x}_i + (t-1)A(\overline{G})\mathbf{x}_i + 2(t-1)\mathbf{x}_i \\ \hline \vdots \\ \hline \overline{q}_i\mathbf{x}_i + ((t-1)(n-1)I_n - (t-1)D)\mathbf{x}_i + (t-1)A(\overline{G})\mathbf{x}_i + 2(t-1)\mathbf{x}_i \end{bmatrix} =$$

$$\begin{bmatrix} \overline{q}_i\mathbf{x}_i + (t-1)D(\overline{G})\mathbf{x}_i + (t-1)A(\overline{G})\mathbf{x}_i + 2(t-1)\mathbf{x}_i \\ \hline \overline{q}_i\mathbf{x}_i + (t-1)D(\overline{G})\mathbf{x}_i + (t-1)A(\overline{G})\mathbf{x}_i + 2(t-1)\mathbf{x}_i \\ \hline \vdots \\ \hline \overline{q}_i\mathbf{x}_i + (t-1)D(\overline{G})\mathbf{x}_i + (t-1)A(\overline{G})\mathbf{x}_i + 2(t-1)\mathbf{x}_i \end{bmatrix} =$$

$$= \begin{bmatrix} \overline{q_i}\mathbf{x}_i + (t-1)Q(\overline{G})\mathbf{x}_i + 2(t-1)\mathbf{x}_i \\ \hline \overline{q_i}\mathbf{x}_i + (t-1)Q(\overline{G})\mathbf{x}_i + 2(t-1)\mathbf{x}_i \\ \hline \vdots \\ \hline \overline{q_i}\mathbf{x}_i + (t-1)Q(\overline{G})\mathbf{x}_i + 2(t-1)\mathbf{x}_i \end{bmatrix} = \begin{bmatrix} t\overline{q}_i\mathbf{x}_i + 2(t-1)\mathbf{x}_i \\ \hline t\overline{q}_i\mathbf{x}_i + 2(t-1)\mathbf{x}_i \\ \hline \vdots \\ \hline t\overline{q}_i\mathbf{x}_i + 2(t-1)\mathbf{x}_i \end{bmatrix} = (t\overline{q}_i + 2(t-1))\mathbf{y}_i.$$

So $\mathbf{y}_i$ is an eigenvector to $t\overline{q}_i + 2(t-1)$. It easy to see that $\mathbf{y}_i^T\mathbf{y}_j = 0$, since $\mathbf{x}_i^T\mathbf{x}_j = 0$, for all $i \neq j$, $1 \leq i, j \leq n$ and the result follows. □

References

[1] Anderson, W.N., and T.D. Morley, *Eigenvalues of the Laplacian of a graph*, Linear and Multilinear Algebra **18** (1985),141–145.